A central data archiving system, said system comprising:

- a data source providing medical data;
- a status monitor for controlling the transfer of said medical data from said data source to a remote data store; and
- a remote data store receiving said medical data and storing said medical data.

[c2]

The system of claim 1, wherein said status monitor verifies said transfer of said medical data from said data source to said/remote data store.

- [c3] The system of claim 1, further comprising an access authenticator for authenticating access to said remote pata store by said data source.
- [c4] The system of claim 3, wherein said access authenticator authenticates access to said data source.
- [c5] The system of claim 1, wherein said data source further stores medical data.
- [c6] The system of claim 5, wherein said remote data store further restores said medical data to said data/source.
 - The system of claim 1/wherein said remote data store stores a copy of said medical data.
- [c8] The system of claim 1, further comprising a second data source for storing medical data, wherein said remote data store transfers said medical data to said second data source.
- [c9] The system of claim 1, wherein said remote data store comprises an application service provider.
- [c10] The system of claim 1, wherein said remote data store is stored at a plurality of locations.
- [c11] The system of claim 1, wherein said status monitor controls the transfer of data from said data source to said remote data store at a definable interval.

[c7]

	[c12]	The system of claim 11, wherein said definable interval comprises a timed
		interval.
	[c13]	The system of claim 11, wherein said definable interval comprises an event-
		based interval.
	[c14]	The system of claim 11, wherein said definable interval comprises a manual
	[614]	interval.
ر	[c15]	A system for remotely accessing a centralized data store, said system
		comprising:
)		a remote data store storing medical data;
		a status monitor for controlling the transfer of said medical data from said
		remote data store to a data source; and
		a data source receiving said medical data and storing said medical data.
	[c16]	The system of claim 15, further comprising a second data source storing
:		medical data.
	[c17]	The system of claim 16, wherein said status monitor controls the transfer of
	[(17]	said copy of said medical data between said remote data store and said second
		data source.
		data source.
	[c18]	The system of claim 16, wherein said status monitor verifies the transfer of said
		copy of said medical data between said remote data store and said second data
		source.
	[c19]	The system of claim 15, further comprising an access authenticator for
	[2.0]	authenticating access to said remote data store.
		· · · · · · · · · · · · · · · · · · ·
	[c20]	The system of claim 15, wherein said status monitor verifies said transfer of
		said medical data between said first data source and said remote data store.
	[c21]	The system of claim 15, wherein said remote data store comprises an
		application service provider.
	[c22]	The system of claim 15, wherein said remote data store is stored at a plurality

	[c23]	The system of claim 15, wherein said remote data store restores said medical
		data at said data source.
	[c24]	The system of claim 15, wherein said remote data store comprises at least one
		directory corresponding to said first data source.
W.	[c25]	A method for remotely archiving medical data, said method comprising:
3		transferring said medical data from a data source to a remote data store; and
/		storing said medical data at said remote data store.
	[c26]	The method of claim 25, further comprising the step of obtaining said medical
		data.
	[c27]	The method of claim 25, further comprising the step of storing said medical
Ď		data at said data source.
#	[c28]	The method of claim 25, wherein said storing step further comprises storing
H		said medical data at said remote data store in a directory corresponding to said
		data source.
<u> </u>	[c29]	The method of claim 25, wherein said transferring step further comprises
		verifying said transfer of medical data from said remote data store to said data
		source.
	[c30]	The method of claim 25, further comprising the step of authenticating access to
		said remote data store.
	[c31]	The method of claim 25, wherein said transferring step occurs after a definable
		interval.
	[c32]	The method of claim 31, wherein said definable interval comprises a timed
		interval.
٠	[c33]	The method of claim 31, wherein said definable interval comprises an event-
		based interval.
		/

of locations.

[c34]	The method of claim 31, wherein said definable interval comprises a manual
	interval.
[c35]	The method of claim 25, further comprising the step of restoring said medical
	data to said data source from said remote data store.
[c36]	The method of claim 25, further comprising the step of copying said medical
	data from said remote data source to a second data source.
[c37]	A method for restoring medical data to a data source from a remote data store,
	said method comprising:
	detecting an error in accessed medical data;
	transferring a copy of said medical data from a remote data store to said data source; and
	restoring said medical data by replacing said medical data at said data source
	with said copy of said medical data.
[c38]	The method of claim 37, further comprising the step of previously obtaining
	said medical data at said data source and storing said medical data at said
	remote data store.
[c39]	The method of claim 37, further comprising the step of copying said medical
	data to a second data source.
[c40]	The method of claim 37, wherein said transferring step further comprises
	verifying said transferring of medical data from said remote data store to said
	data source.
[c41]	The method of claim 37, further comprising the step of authenticating access to
	said remote data store.
[c42]	The method of claim 37, wherein said transferring step further comprises
	transferring said medical data from a directory representative of said data
	source at said remote data store to said data source.
[c43]	
	A method for installing medical data from a first data source to a second data

[c44]

[c45]

[c46]

[c47]


[c48]

[c49]

[c50]

[c51]

[c52]



The method of claim 49, wherein said definable interval comprises a manual

interval.

This Page Blank (uspto)